

ON A CLASS OF WEAKLY BERWALD SPECIAL (A, B) -METRICS OF SCALAR FLAG CURVATURE

Vanithalakshmi S M^{1*} and Narasimhamurthy S K², Roopa. M. K³

^{*123}Department of Mathematics, Kuvempu University, Shankaraghatta, Shimoga, Karnataka, India

²e-mail: nmurthysk@gmail.com and ³Email: roopamk1002@gmail.com

***Corresponding Author:-**

Email: vanithateju02@gmail.com

Abstract:-

In this paper, we study the Finsler space with special (α, β) -metric $F = \alpha + \frac{\beta^2}{\alpha}$ is scalar flag curvature and we proved that, if it is weakly Berwald if and only if it is Berwald and vanishes flag curvature. Further, we found that this metric is locally Minkowskian.

2010 AMS Subject Classification: 53C25, 53C40, 53D15.

Key Words:-Finsler space, S-curvature, (α, β) -metrics, Mean Berwald curvature, Homogenous Finsler space.

1. INTRODUCTION

In Finsler geometry, the flag curvature is an analogue of the sectional curvature in Riemannian geometry. One of the fundamental problem in Riemannian-Finsler geometry is to study and characterize Finsler metrics of scalar flag curvature. It is known that every Berwald metric is a Landsberg metric and also, for any Berwald metric the S -curvature vanishes. In 2003 X.Chang, X.Mo and Shen have obtained the results on the flag curvature of Finsler metrics of scalar curvature. Then, in 2004, Yoshikawa, Okubo and M. Matsumoto showed the conditions for some (α, β) -metrics to be weakly Berwald. Recently Narasimhamurthy.S.K[8] has studied the class of weakly Berwald (α, β) -metrics of scalar flag curvature.

The main purpose of the present paper is to characterize the weakly Berwald special (α, β) -metrics of scalar flag curvature. Then, we proved that it vanishes scalar flag curvature and Berwald, then it is locally Minkowskian.

2. Preliminaries:

This section includes the basics of Finsler spaces and the concepts of S -curvature.

Let $F = \alpha \phi \left(\frac{\beta}{\alpha} \right)$ be an (α, β) -metric on an n -dimensional manifold M , where $\alpha = \sqrt{a_{ij}(x)y^i y^j}$ is a Riemannian metric and $\beta = b_i(x)y^i$ is a 1-form.

Now, let $\phi(s)$ be a positive c^∞ function on $(-b_0, b_0)$. For a number $b \in [0, b_0)$, let

$$\phi = -(Q - sQ')n\Delta + 1 + sQ - (b^2 - s^2)(1 + sQ)Q'', \quad (2.1)$$

where, $\Delta = 1 + sQ + (b^2 - s^2)Q'$ and $Q = \frac{Q}{\phi - sQ'}$.

Let G^i and \tilde{G}^i denotes the spray co-efficients of F and α respectively. Then geodesic co-efficients G^i are related to \tilde{G}^i of by

$$G^i = \tilde{G}^i + \alpha Q s_0^i + \Theta \{-2\alpha Q s_0 + r_{00}\} \left\{ \frac{y^i}{\alpha} + \frac{Q'}{Q - sQ'} b^i \right\}, \quad (2.2)$$

where \tilde{G}^i denote the spray coefficients of α .

We shall denote $r_{00} = r_{ij}y^i y^j$, $s_{i0} = s_0 = s_i y^i$ and $\Theta = \frac{Q - sQ'}{2\Delta}$.

Also,

$$h_i = \alpha b_i - s y_i, \quad \psi_i = \sqrt{b^2 - s^2} \Delta^{1/2} \left[\frac{\sqrt{b^2 - s^2}}{\Delta^{3/2}} \Phi \right]',$$

$$\psi_2 = 2(n+1)(Q - sQ') + 3 \frac{\phi}{\Delta}.$$

By a direct computation, we can obtain a formula for the mean Cartan torsion of (α, β) -metrics as follows[5].:

$$I_i = -\frac{\Phi(\phi - s\phi')}{2\Delta\phi\alpha^2} (\alpha b_i - s y_i). \quad (2.3)$$

According to Diecke's theorem[10], a Finsler metric is Riemannian if and only if the mean Cartan torsion vanishes, $I = 0$.

Clearly, an (α, β) -metric $F = \alpha\phi(\beta/\alpha)$, $s = \beta/\alpha$ is Riemannian if and only if $\Phi = 0$. (see [21]).

For a Finsler metric $F = F(x, y)$ on a manifold M , the Riemann curvature $R_y = R_k^i \frac{\partial}{\partial x^i} \otimes dx^k$ is defined by

$$R_k^i = 2 \frac{\partial G^i}{\partial x^k} - \frac{\partial^2 G^i}{\partial x^m \partial y^k} y^m + 2G^m \frac{\partial^2 G^i}{\partial y^m \partial y^k} - \frac{\partial G^i}{\partial y^m} \frac{\partial G^m}{\partial y^k}.$$

Let $R_{jk} = g_{ji} R_{ki}$, then $R_{jkyj} = 0$, $R_{jk} = R_{kj}$.

For a flag $\{P, y\}$, where $P = \text{span}\{y, u\} \subset T_x M$, the flag curvature $K = K(P, y)$ of F is defined by

$$K(P, y) = \frac{R_{jk}(x, y) u^j u^k}{F^2(x, y) h_{jk}(x, y) u^j u^k}$$

$$h_{jk} = g_{jk} - F^{-2} g_{jp} y^p g_{kq} y^q. \quad (2.4)$$

where,

We say that Finsler metric F is of scalar flag curvature if the flag curvature $K = K(x, y)$ is independent of the flag P . By the definition, F is of scalar flag curvature $K = K(x, y)$ if and only if in a standard local coordinate system,

$$R_k^i = KF^2 h^i_k, \quad (2.5)$$

where, $h^i_k = g^{ij} h_{jk} = g^{ij} F F_{y^j k}$ ([4], [5]).

The Schur lemma[4] in Finsler geometry tell us that, in dimension $n \geq 3$, if F is of isotropic flag curvature, $K = K(x)$, then it is of constant flag curvature, $K = \text{constant}$. The Berwald curvature $B_y = B_{jkl}^i dx^j \otimes \frac{\partial}{\partial x^i} \otimes dx^k \otimes dx^l$ and mean Berwald curvature $E_y = E_{ij} dx^i \otimes dx^j$ are defined respectively by

$$B_{jkl}^i = \frac{\partial^3 G^i(x, y)}{\partial y^j \partial y^k \partial y^l}, \quad E_{ij} = \frac{1}{2} \frac{\partial^2}{\partial y^i \partial y^j} \left(\frac{\partial G^m}{\partial y^m} \right) = \frac{1}{2} B_{ij}^m.$$

A Finsler metric F is called weak Berwald metric if the mean Berwald curvature vanishes, i.e., $(E = 0)B = 0$. A Finsler metric F is said to be of isotropic mean Berwald curvature if $F = \frac{1}{2}(n + 1)c(x)F^{-1}h$, where $c = c(x)$ is a scalar function on the manifold M .

Theorem 2.1. [8],[9] For special (α, β) -metric $F = \alpha + k\frac{\beta^2}{\alpha}$, where $k \neq 0$, is constant and Matsumoto metric (2nd appx) $F = \alpha + \beta + \frac{\beta^2}{\alpha} + \frac{\beta^3}{\alpha^2}$ on n -dimensional manifold M . Then the following are equivalent:

- (a) F is of isotropic S -curvature, $S = (n + 1)c(x)F$;
- (b) F is of isotropic mean Berwald curvature, $E = \frac{n+1}{2}c(x)F^{-1}h$;
- (c) β is a killing 1-form with constant length with respect to α , i.e., $r_{00} = 0$ and $s_{00} = 0$;
- (d) S -curvature vanishes, $S = 0$;
- (e) F is a weak Berwald metric $E = 0$; where, $c = c(x)$ is scalar function on the manifold M .

Note that, the discussion in [3], [19] doesn't involve whether or not F is Berwald metric. By the definition [7], Berwald metrics must be weak Berwald metrics but the converse is not true. For this observation, we further study the Finsler metrics $F = \alpha + \frac{\beta^2}{\alpha}$.

For a Finsler metric $F = F(x, y)$ on an n -dimensional manifold M , the Busemann-Hausdorff volume form $dV_F = \sigma_F(x)dx^1 \wedge \dots \wedge dx^n$ is given by

$$\sigma_F(x) = \frac{\text{vol}(B^n(1))}{\text{vol}\{(y^i) \in R^n | F(x, y) < 1\}},$$

Vol denotes the Euclidean volume in R^n . The S -curvature is given by

$$S(x, y) = \frac{\partial G^m}{\partial y^m} - y^m \frac{\partial \ln \sigma_F}{\partial x^m}. \tag{2.6}$$

Clearly, the mean Berwald curvature $E_y = E_{ij}dx^i \otimes dx^j$ can be characterized by use of S -curvature as follows:

$$E_{ij} = \frac{1}{2} \frac{\partial^2 S}{\partial y^i \partial y^j},$$

A Finsler metric F is said to be isotropic S -curvature if $S = (n+1)c(x)F(x, y)$, where $c = c(x)$ is a scalar function on the manifold M . S -curvature is closely related to the flag curvature. We use the following important results proved by [7].

Theorem 2.2. Let (M, F) be an n -dimensional Finsler manifold of scalar flag curvature with flag curvature $K = K(x, y)$. Suppose that the S -curvature is isotropic, $S = (n+1)c(x)F(x, y)$, where $c = c(x)$ is a scalar function on M . Then there is a scalar function $\sigma(x)$ on M such that

$$K = \frac{3c_{x^m}(x)y^m}{F(x, y)} + \sigma(x). \tag{2.7}$$

This shows that S -curvature has important influence on the geometric structures of Finsler metrics. For a Finsler metric F , the Landsberg curvature $L = L_{ijk}dx^i \otimes dx^j \otimes dx^k$ and the mean Landsberg curvature $L = J_k dx^k$ are defined respectively by

$$L_{ijk} = -\frac{1}{2} F F_{y^m} [G^m]_{y^i y^j y^k}, \quad J_k = g^{ij} L_{ijk},$$

A Finsler metric F is called weak Landsberg curvature metric if the mean Landsberg curvature vanishes, i.e., $(J = 0)L = 0$. For an (α, β) -metric $F = \alpha\phi(s)$, $s = \frac{\beta}{\alpha}$, Li Benling and Z, Shen [7]

obtained the following formula of the mean Landsberg curvature

$$J_i = -\frac{1}{2\Delta\alpha^4} \left\{ \frac{2\alpha^2}{b^2 - s^2} \left[\frac{\phi}{\Delta} + (n + 1)(Q - sQ') \right] (s_0 + r_0)h_i + \frac{\alpha}{b^2 - s^2} \left[\psi_1 + s\frac{\phi}{\Delta} \right] (r_{00} - 2Q\alpha s_0)h_i \right. \\ \left. + \alpha \left[-\alpha Q' s_0 h_i + \alpha Q(\alpha^2 s_i - y_i s_0) \right] + \alpha^2 \Delta s_{i0} + \left[\alpha^2 (r_{i0} - 2\alpha Q s_i) - (r_{00} - 2\alpha Q s_0) y_i \right] \frac{\phi}{\Delta} \right\}. \tag{2.8}$$

Besides, they also obtained

$$\bar{J} = J_i b^i = -\frac{1}{2\Delta\alpha^2} \left\{ \psi_1 (r_{00} - 2\alpha Q s_0) + \alpha \psi_2 (r_0 + s_0) \right\}. \tag{2.9}$$

The horizontal covariant derivatives $J_{i;m}$ and $J_{i|_m}$ of J_i with respect to F and α respectively are given by

$$J_{i;m} = \frac{\partial J_i}{\partial x^m} - J_l \Gamma_{im}^l - \frac{\partial J_i}{\partial y^l} N_m^l, \quad J_{i|m} = \frac{\partial J_i}{\partial x^m} - J_l \bar{\Gamma}_{im}^l - \frac{\partial J_i}{\partial y^l} \bar{N}_m^l,$$

where,

$$\Gamma_{ij}^l = \frac{\partial G^l}{\partial y^i \partial y^j}, \quad N_j^l = \frac{\partial G^l}{\partial y^j} \quad \text{and} \quad \bar{\Gamma}_{ij}^l = \frac{\partial \bar{G}^l}{\partial y^i \partial y^j}, \quad \bar{N}_j^l = \frac{\partial \bar{G}^l}{\partial y^j}.$$

Further we have,

$$\begin{aligned} J_{i;m} y^m &= \{J_{i|m} - J_l (\Gamma_{im}^l - \bar{\Gamma}_{im}^l) - \frac{\partial J_i}{\partial y^l} (N_m^l - \bar{N}_m^l)\} y^m \\ &= J_{i|m} y^m - J_l (N_i^l - \bar{N}_i^l) - 2 \frac{\partial J_i}{\partial y^l} (G^l - \bar{G}^l). \end{aligned}$$

If a Finsler metric F is of constant flag curvature K (see[5]), then

$$J_{i;m} y^m + KF2I_i = 0.$$

So, if can (α, β) -metric $F = \alpha \phi(s)$, $s = \frac{\beta}{\alpha}$, is of constant flag curvature K , then

$$J_{i|m} y^m - J_l \frac{\partial (G^l - \bar{G}^l)}{\partial y^i} - 2 \frac{\partial J_i}{\partial y^i} (G^l - \bar{G}^l) + K \alpha^2 \phi^2 I_i = 0.$$

Contracting the above equation by b^i yields the following equation

$$\bar{J}_{i|m} y^m - J_i \alpha^{ik} b_{k|m} y^m - J_l \frac{\partial (G^l - \bar{G}^l)}{\partial y^i} b^i - 2 \frac{\partial \bar{J}}{\partial y^l} (G^l - \bar{G}^l) + K \alpha^2 \phi^2 I_i b^i = 0 \quad (2.10)$$

3. Characterization of Weakly Berwald special (α, β) -metrics of scalar flag curvature:

In this section, for a n -dimensional Finsler manifold ($n \geq 3$), we characterize the class of weakly Berwald special (α, β) -metric of scalar flag curvature. So first we prove the following lemma;

Lemma 3.1. Let $F = \alpha + \frac{\beta^2}{\alpha}$ be a Finsler space with (α, β) -metric, then it is non-Randers type, if $\Theta = 0$.

Proof: Consider the Finsler metric $F = \alpha + \frac{\beta^2}{\alpha}$. By direct computation using (2.1) we have

$$\Theta = \frac{-A\phi}{(1-s^2)^4},$$

where,

$$\phi = 1 + s^2$$

$$A = (3n + 1)s^5 - 16s^4 - (4(n + 1) + 8nb^2 + 4b^2) s^3 + 4(4b^2 - 1)s^2 + 4b^2.$$

Assume that $\Theta = 0$, then $A = 0$ obviously and so multiplying $A = 0$ with α^5 , which yields.

$4\beta^5(1 + 12n) - [(n + 1) + 2nb^2 + B^2]\beta^3\alpha^2 + \alpha[4b^2\alpha^4 + 4(4b^2 - 1)\beta^2\alpha^2 - 16\beta^4] = 0$. Hence we obtain,

$$\beta^5(1 + 12n) - ((n + 1) + (2n + 1)b^2) \beta^3 \alpha^2 = 0, \quad (3.1)$$

and

$$b^2\alpha^4 + (4b^2 - 1)\beta^2\alpha^2 - 4\beta^4 = 0. \quad (3.2)$$

Clearly, observe that in equation (3.1) β^5 is divisible by α^2 , which is contraction to $\Theta = 0$.

Therefore, F must be a non-Randers type.

The main idea of this lemma(3.1), now we can prove the following theorem.

Theorem 3.3. Let $F = \alpha + \frac{\beta^2}{\alpha}$ be a Finsler space with (α, β) -metric is of scalar flag curvature $K = K(x, y)$, then F is weakly Berwald metric if and only if F is Berwald metric and $K = 0$.

In this case, F must be locally Minkowskian.

Proof. From lemma(3.1) and theorem(2.2), we know that Finsler space with (α, β) -metrics $F = \alpha + \frac{\beta^2}{\alpha}$ cannot represents the Riemannian mertric if $\beta \neq 0$.

Then the proof of this theorem sufficient part is trivial. and so we only prove the necessary condition.

Assume that the metric F is weakly Berwald. By lemma(3.1) and the isotropic S -curvature, $S = (n + 1)c(x)F$ with $c(x)$ is constant and

$$r_{00} = 0, \quad s_0 = 0. \quad (3.3)$$

Again theorem(2.2) and the equation (2.7) F must be of isotropic flag curvature $K = \sigma(x)$.

Further, equation(2.10), we can simply using (2.2),(2.8) and (2.9) as follows.

$$G^i - \bar{G}^i = \alpha Q s_0^i, \quad J_i = \frac{-\Theta s_{i0}}{2\alpha\Delta}, \quad \bar{J} = 0.$$

In addition from(2.3) we obtain

$$I_i b^i = -\frac{\Theta(\phi - s\phi')}{2\Delta F}(b^2 - s^2), \quad (3.4)$$

thus, equation(2.10) can be expressed as follows:

$$\frac{\Theta s_{i0}}{2\Delta\alpha} a^{ik} s_{k0} + \frac{\Theta s_{i0}}{2\Delta\alpha} (sQs_0^l + Q's_0^l(b^2 - s^2)) - KF\frac{\Theta}{2\Delta}(\phi - s\phi')(b^2 - s^2) = 0.$$

Again by using lemma(3.1) we get

$$s_{i0}s_0^i + s_{i0}(\alpha Qs_0^l)b_i - KF\alpha(\phi - s\phi')(b^2 - s^2) = 0.$$

Here, notice that, $F = \alpha\phi(s)$, $s = \frac{\beta}{\alpha}$. Thus we have,

$$s_{i0}s_0^i\Delta - K\alpha^2\phi(\phi - s\phi')(b^2 - s^2) = 0. \quad (3.5)$$

For Finsler space with (α, β) -metric $F = \alpha + \frac{\beta^2}{\alpha}$ we have,

$$\Delta = \frac{\phi(1+2b^2-3s^2)}{(1-s^2)^2},$$

Then equation (3.5) becomes,

$$(1 + 2b^2 - 3s^2)s_{i0}s_0^i - K\alpha^2(b^2 - s^2)(1 - s^2)^3 = 0.$$

On multiplying above equation with α^6 , which yields.

$$-b^2\alpha^8 + \{\beta^2(1+3b^2) + (1+2b^2)s_{i0}s_0^i\}\alpha^6 - 3\beta^2\{(1+2b^2) + s_{i0}s_0^i\}\alpha^4 + K\beta^6(3+b^2)\alpha^2 = K\beta^8. \quad (3.6)$$

Here, we observe that, the leftside of equation (3.6) is divisible by α^2 . Hence we obtain that the flag curvature $K = 0$, because β^8 is not divisible by α^2 . Substituting $K = 0$ into equation (3.5) we have $s_{i0}s_0^i = a_{ij}(x)s_0^j s_0^i = 0$. Because $(a_{ij}(x))$ is positive definite, so $s_0^i = 0$, i.e., β is closed. By equation (3.3), we know that β is parallel with respect to α . Then F is Berwald metric if $\beta = 0$. Hence, F must be locally Minkowskian.

References

- [1]. Cheng.X, *On (α, β) -metrics of scalar flag curvature with constant S-curvature*. Acta Math, sin. Engl. ser 26(9), 1701-1708(2010).
- [2]. Cui.N.W, *On the S-Curvature of some (α, β) -metrics*, Acta Math Sci 26A(7), 1047-1056.
- [3]. C.Nigwei, *On the S-Curvature of (α, β) -metrics*, Acta Math Sci 26A(7), 1047-1056.
- [4]. D.Bao, S.S.Chen and Z.Shen, *An Introduction to Riemannian-Finsler Geometry*, Springer Verlag 2000.
- [5]. S.S.Chern and Z.Shen, *Riemannian-Finsler geometry*, World-Scientific publishing Co., Singapore .
- [6]. Laurian-Ioan piscoran. Vishnu Narayan Mishra, *S-Curvature for a new class of (α, β) -metrics*, Splinger-Verlag Italia (2016).
- [7]. Li Benling and Z.Shen, *On a Class of weak Landsberg metrics*, J.Science China Series., 573-589, 50(2004).
- [8]. Narasimhamurthy.S.K, Kavyashree A.R, Ajith and Mallikarjun.Y.K, *On class of weakly Berwald (α, β) metrics of scalar flag Curvature*. , Int. J.Math.Res.5(1), 97-108 (2013).
- [9]. Narasimhamurthy.S.K, Thippeswamy.K.R, *Two kinds of weakly Berwald special (α, β) -metrics of scalar flag Curvature*. , Int. J.of Mathematics Trends and Technology. Vol.44, no-3 (2013).
- [10]. P.L.Antonelli, R.S.Ingarden and M.Matsumoto, *The theory of sprays and Finsler spaces with applications in Physics and biology*. , Kluwer Acad., Dordecht.
- [11]. S.Deng and X.Wang , *S-Curvature of homogeneous (α, β) -metrics*, Balkan Journal of Geometry and its application. Vol.15 ,No.2, pp 39-48 (2010).
- [12]. S.Deng and .Hou, *Homogeneous Finsler spaces of negative curvature School of Mathematical sciences and LPMC, Nankai University, Tianjin, 300071, PR China.*
- [13]. Shaoqing Deng,, *The S-Curvature of homogeneous Randers spaces*, School of Mathematical Sciences, Nankai University, Tianjin 300071, PR China.
- [14]. S.C.Shibata , *On Finsler spaces with Kropina metric*, Rep. Math, Phys. **13** (1978), 117-128.
- [15]. S.S. Chern and Z.Shen, *Riemann Finsler Geometry*, World Scientific Publishers (2004).
- [16]. S.Deng and X,Wang *The S-Curvature of homogeneous (α, β) -metrics* Balkan Journal of Geometry and its applications, Vol 15, No.2, pp. 39-48, (2010).
- [17]. S.Bacso, X.Cheng and Z.Shen, *Curvature properties of (α, β) -metrics*, Adv. study, pure Math. Sco., Japan(2007).
- [18]. V.K. Kropina, *On Projective Finsler space with a certain special form*, Nauch Doklady vyss. Skoly, fiz-math Nauki 19-59(2), 38-42, Russian(1960).
- [19]. Vanithalakshmi.S.M and Narasimhamurthy.S.K, *S- Curvature of homogeneous Finsler space with special (α, β) -metrics*, Gulf Journal of Mathematics, 1-11 vol 1(2018).
- [20]. X.Cheng and Z.Shen, *A Class of Finsler metrics with isotropic S-Curvature*, Isreal J. Math 169, (2009).
- [21]. X.Cheng and H.Wang and M.Wang, *(α, β) -metrics with relatively isotropic mean Landsberg curvaturre*, J.Publ.Math. Debraceen, 475-485, 72(2008) 169.
- [22]. Z.Hu and S.Deng, *Three dimensionssal homogeneous Finsler manifolds*, Math. Nachr. 285, No. 10, 12431254(2012).
- [23]. Z.Shen *Volume comparision and its applications in Riemannian Finsler geometry* , Adv.Math. 128, 306-328, (1997).